

Increase in intranuclear nuclear factor κ B and decrease in inhibitor κ B in mononuclear cells after a mixed meal: evidence for a proinflammatory effect¹⁻³

Ahmad Aljada, Priya Mohanty, Husam Ghanim, Toufic Abdo, Devjit Tripathy, Ajay Chaudhuri, and Paresh Dandona

ABSTRACT

Background: In view of the stimulatory effect of glucose on reactive oxygen species (ROS) generation, we investigated the possibility that a mixed meal stimulates ROS generation and possibly induces concomitant proinflammatory changes.

Objective: The objective was to determine whether the intake of a 900-kcal mixed meal induces an increase in ROS generation by leukocytes and an inflammatory response at the cellular level.

Design: Nine normal-weight subjects were given a 900-kcal mixed meal, and 8 normal-weight subjects were given 300 mL water after an overnight fast. Blood samples were collected at 0, 1, 2, and 3 h. ROS generation by mononuclear cells and polymorphonuclear leukocytes and the expression of p47^{phox} subunit were measured. Intranuclear nuclear factor κ B (NF- κ B) binding and the expression of inhibitor κ B α (I κ B α), I κ B kinase α (IKK α), and I κ B kinase β (IKK β) were measured. Plasma concentrations of C-reactive protein (CRP) and soluble intercellular adhesion molecule were also measured.

Results: ROS generation by mononuclear cells and polymorphonuclear leukocytes and p47^{phox} expression increased significantly. The expression of IKK α and IKK β and DNA-binding activity of NF- κ B increased significantly, whereas I κ B α expression decreased. Plasma CRP concentrations increased. The intake of 300 mL water did not induce a change in any of the above indexes.

Conclusions: These data show that the intake of a mixed meal results in significant inflammatory changes characterized by a decrease in I κ B α and an increase in NF- κ B binding, plasma CRP, and the expression of IKK α , IKK β , and p47^{phox} subunit. These proinflammatory changes are probably relevant to the state of chronic hypertension and obesity and to its association with atherosclerosis. *Am J Clin Nutr* 2004;79:682-90.

KEY WORDS Nuclear transcription factor κ B, inhibitor κ B, mononuclear cells, mixed meal, inflammation, reactive oxygen species

INTRODUCTION

We recently showed that glucose (1) and lipid and protein (2) intakes cause an increase in the generation of reactive oxygen species (ROS) by leukocytes and that each of these macronutrients induces a distinct pattern of increase in ROS generation. Thus, glucose induces a peak in ROS generation by both mononuclear cells (MNCs) and polymorphonuclear leukocytes

(PMNLs) at 2 h, whereas cream (lipid) produces a peak at 1 h. The peak increase in ROS generation is the greatest with glucose, whereas it is the least with casein (protein). On the other hand, cream intake causes a prolonged increase in lipid peroxidation. Consistent with these observations, we also showed that a 48-h fast results in a marked decrease in ROS generation by leukocytes and oxidative damage of amino acids (3). Thus, we suggest that nutritional intake may be the major modulator of ROS generation. Indeed, we showed recently that the state of obesity reflecting chronic hypernutrition is associated with marked oxidative stress, as reflected in an increase in indexes of lipid peroxidation, protein carbonylation, and oxidative damage of amino acids, which diminishes after dietary restriction over a relatively short period of time, eg, 4 wk (4).

Because our studies to date have not dealt with real food, but instead with pure macronutrients, we have now undertaken our first study with food items. We chose to study the effect of a mixed meal composed of carbohydrates, protein, and fats. Such a mix was previously shown to impair endothelium-mediated vasodilatation in the brachial artery (5, 6). In addition, Ceriello et al (7) showed an increase in oxidative stress and LDL oxidation in diabetes after a meal challenge.

We hypothesized that the intake of a mixed meal causes an increase in oxidative stress and proinflammatory activity in circulating MNCs. In this study we investigated the effect of the intake of this mixed meal on ROS generation by PMNLs and MNCs. Because we previously showed that glucose intake results in an increase in the p47^{phox} subunit, the key protein of NADPH oxidase in leukocytes, an increase in intranuclear nuclear factor κ B (NF- κ B), and a decrease in I κ B α (8), we decided to also investigate the effect of the mixed meal on intranuclear NF- κ B binding activity and the expression of inhibitor κ B α (I κ B α), I κ B kinase α (IKK α), I κ B kinase β (α IKK β), and p47^{phox} subunit in MNC. Plasma C-reactive protein (CRP), soluble intercellular adhesion molecule 1 (sICAM-1), and α -

¹ From the Division of Endocrinology, Diabetes and Metabolism, State University of New York at Buffalo and Kaleida Health, Buffalo, NY.

² Supported by the William G McGowan Charitable Fund, Washington, DC.

³ Address reprint requests to P Dandona, Diabetes-Endocrinology Center of Western New York, 3 Gates Circle, Buffalo, NY 14209. E-mail: pdandona@kaleidahealth.org.

Received November 27, 2002.

Accepted for publication September 16, 2003.

TABLE 1

Glucose, insulin, triacylglycerol, α -tocopherol, and total, HDL-, and LDL-cholesterol concentrations after a 900-kcal mixed-meal challenge¹

	Time after the meal challenge			
	Baseline (0 h)	1 h	2 h	3 h
Glucose (mg/dL)	82 \pm 3 ²	95 \pm 6	89 \pm 6	93 \pm 3
Insulin (μ U/mL) ³	6.1 (5.7–6.6)	28.8 (22.7–72.6) ⁴	17.5 (14.1–32.0) ⁴	19.8 (13.4–14.6) ⁴
Triacylglycerol (mg/dL)	97 \pm 14	121 \pm 13	146 \pm 17 ⁴	173 \pm 20 ⁴
HDL (mg/dL)	57 \pm 14	54 \pm 12	52 \pm 10	53 \pm 12
LDL (mg/dL)	104 \pm 8	90 \pm 7	85 \pm 8	81 \pm 11
Cholesterol (mg/dL)	180 \pm 12	168 \pm 11	165 \pm 9	169 \pm 12
α -Tocopherol (μ g/mL)	15.3 \pm 2.5	15.2 \pm 2.5	14.6 \pm 2.3	14.7 \pm 1.8

¹ $n = 9$.² $\bar{x} \pm$ SEM (all such values).³ Median; interquartile range in parentheses.⁴ Significantly different from baseline, $P < 0.05$ (one-factor ANOVA for repeated measures followed by Dunnett's test).

tocopherol were also measured. These indexes would allow us to explore a possible link between nutrition, oxidative stress, and inflammation.

SUBJECTS AND METHODS

Subjects

Nine nondiabetic subjects aged 29–38 y ($\bar{x} \pm$ SEM: 32 \pm 3 y) of normal weight [body mass index (in kg/m²): 25.4 \pm 2.7; $\bar{x} \pm$ SEM] were included in the study. The subjects came to the Clinical Research Center of the Diabetes–Endocrinology Center of Western New York after an overnight fast between 0800 and 0900. A fasting blood sample was obtained, and the subjects were asked to eat a mixed meal containing 910 kcal (egg-muffin and sausage-muffin sandwiches and 2 hash browns, which contained 81 g carbohydrate, 51 g fat, and 32 g protein) over 15 min. Additional blood samples were obtained 1, 2, and 3 h after the meal was eaten. The control subjects (age: range, 26–50 y; $\bar{x} \pm$ SD, 35 \pm 8 y; body mass index: 24.3 \pm 2.0) were 8 volunteers who were given 300 mL water to drink in the fasting state. Blood samples were obtained before and 1, 2, and 3 h after the drink. The protocol was approved by the Institutional Review Board of the State University of New York at Buffalo based at The Millard Fillmore Hospital. All participants gave their written informed consent.

Isolation of PMNLs and MNCs

Blood samples were collected in tubes containing Na-EDTA as an anticoagulant; 3.5 mL of the anticoagulated blood sample was carefully layered over 3.5 mL of the PMNL isolation medium (Robbins Scientific Corp, Sunnyvale, CA). Samples were centrifuged at 450 \times g in a swing outrotor for 30 min at 22 °C. At the end of centrifugation, 2 bands separate out at the top of the red blood cell pellet. The top band consists of MNCs, whereas the bottom band consists of PMNLs. The MNC and PMNL bands were harvested with a Pasteur pipette, repeatedly washed with Hank's balanced salt solution, and reconstituted to a concentration of 4 \times 10⁵ cells/mL in Hank's balanced salt solution.

NF- κ B electrophoretic mobility shift assay

DNA-binding protein extracts were prepared from MNCs with the method described by Andrews and Faller (9). Total protein concentrations were determined with the bicinchoninic

acid protein assay (Pierce, Rockland, IL). The NF- κ B gel retardation assay was performed with the use of a NF κ B binding protein detection kit (Life Technologies, Long Island, NY). Briefly, the double-stranded oligonucleotide containing a tandem repeat of the consensus sequence for the NF- κ B binding site was radiolabeled with γ -P³² by T4 kinase. Then, 5 μ g of the nuclear extract was mixed with the incubation buffer, and the mixture was preincubated at 4 °C for 15 min. Labeled oligonucleotide (60 000 cpm) was added, and the mixture was incubated at room temperature for 20 min. Samples were then applied to wells of 6% nondenaturing polyacrylamide gel. The gel was dried under vacuum and exposed to X-ray film. Densitometry was performed with the use of Bio-Rad molecular analyst software (Hercules, CA).

p47^{phox} Subunit, I κ B α , IKK α , and IKK β Western blotting

Western blotting was carried out as previously described. Briefly, total protein concentrations were determined with the use of a bicinchoninic acid protein assay (Pierce, Rockland, IL); 20 μ g MNC homogenate was electrophoresed on sodium dodecyl sulfate polyacrylamide gels for the p47^{phox} subunit. Forty micrograms of total homogenates was used for I κ B α (Cell Signaling Technology Inc, Beverly, MA), IKK α (PharMingen, San Diego), and IKK β (Transduction Labs, Lexington, KY) Western blotting.

Measurement of reactive oxygen species generation

Five hundred microliters of PMNLs or MNCs (2 \times 10⁵ cells) were delivered into a Chronolog Lumi-Aggregometer cuvette. Fifteen microliters of 10 mmol luminol/L was then added, followed by 1.0 μ L of 10 mmol formylmethionyl leucyl phenylalanine/L. Chemiluminescence was recorded for 15 min (a protracted record after 15 min did not alter the relative amounts of chemiluminescence produced by various blood samples). Our method, developed independently (10, 11), is similar to that published by Tosi and Hamedani (12). In this assay system, the release of superoxide radical as measured by chemiluminescence has been shown to be linearly correlated with that measured by the ferricytochrome C method (12). We further established that, in our assay system, there is a dose-dependent inhibition of chemiluminescence by superoxide dismutase and catalase as well as diphenylene iodonium (data not shown), which is a spe-




FIGURE 1. A: Representative electrophoretic mobility shift assay showing the relative nuclear transcription factor κ B (NF- κ B) binding to the double-stranded oligonucleotide containing the NF- κ B DNA-binding site after a 900-kcal mixed-meal challenge or water intake. Band-shift assays were performed with the use of 5 μ g mononuclear cell extract for each time point. Lane 1 represents 0 h, lane 2 represents 1 h, lane 3 represents 2 h, and lane 4 represents 3 h after meal intake. Two complexes were detected in the gels (I and II). The super shift assay showed that complex I is composed of p50:p65 subunits, whereas complex II is composed of p50:p50 subunits. Lane 5: super shift assay with the use of an antibody against the p65 subunit. Lane 6: super shift assay with the use of an antibody against the p50 subunit. Lane 7: antibody against *c-fos* showing no effect with the electrophoretic mobility shift assay. Lane 8: inhibition of NF- κ B binding with the use of 2 μ L competitor NF- κ B oligonucleotide. Lane 9: nonspecific oligonucleotide (AP-1) that shows no effect on binding. This gel clearly shows an increase in complex I formation after the 900-kcal meal challenge and a decrease in the p50:p50 complex (II); however, the decrease in the p50:p50 complex was not consistent in other subjects and did not change significantly (data not shown). B: Mean (\pm SEM) change in NF- κ B binding activity for complex I after intake of the mixed meal by densitometric quantitative analysis. There was a significant interaction between treatment (water compared with meal) and time ($P < 0.005$, two-factor ANOVA). *Significantly different from baseline (0 h), $P < 0.05$ (one-factor ANOVA for repeated measures followed by Dunnett's test). $n = 9$ for meal intake and $n = 8$ for water intake. C: Area under the curve (AUC) for NF- κ B binding activity after water and meal intakes. *Significantly different from water intake, $P < 0.05$ (Student's *t* test).

cific inhibitor of NADPH oxidase—the enzyme responsible for the production of superoxide radicals. The specific inhibitory effect of diphenylene iodonium on NADPH oxidase was established by Hancock and Jones (13).

Measurement of plasma vitamin E, sICAM-1, CRP, and insulin

α -Tocopherol was measured in plasma by HPLC as described previously (1, 4). Plasma sICAM-1 was assayed with an enzyme-linked immunosorbent assay kit from R&D systems (Minneapolis). The intraassay CV for sICAM-1 is 4.6%, whereas the interassay CV is 6.1%. The CRP (intraassay CV: 4%; interassay CV: 5%) and insulin (intraassay CV: 2.6%; interassay CV: 6.2%)

enzyme-linked immunosorbent assay kits were purchased from Diagnostic Systems Laboratories Inc (Webster, TX).

Statistical analysis

The statistical analysis was carried out by using SIGMASTAT software (version 2.03; Jandel Scientific, San Rafael, CA). The analysis was carried out with one-factor repeated-measures analysis of variance (ANOVA) with the use of Dunnett's test for comparisons against the baseline (0 h) for normally distributed data. Dunn's test was used for the nonparametric data. Two-factor ANOVA was used to evaluate the interaction between treatment (water compared with the mixed meal) and time. Student's *t* test was used to compare the area under the curves




FIGURE 2. A: Representative Western blot showing the relative expression of inhibitor κ B α (I κ B α) in mononuclear cell homogenates 0, 1, 2, and 3 h after a 900-kcal mixed-meal challenge or water intake. B: Mean (\pm SEM) change in I κ B α protein concentrations in mononuclear cells by densitometric quantitative analysis. There was a significant interaction between treatment (water compared with meal) and time ($P < 0.05$, two-factor ANOVA). *Significantly different from baseline (0 h), $P < 0.05$ (one-factor ANOVA for repeated measures followed by Dunnett's test). $n = 9$ for meal intake and $n = 8$ for water intake. C: Area under the curve (AUC) for I κ B α protein expression after water and meal intakes. *Significantly different from water intake, $P < 0.05$ (Student's t test).

(AUCs). For the comparisons of ROS generation by PMNLs and MNCs before and after the meal challenge, the data were normalized to a baseline value of 100% because the baseline ROS generation varied markedly from one person to another. We previously showed this same level of variability while showing a consistency (reproducibility) in ROS generation within individuals (10). The AUC for all ROS generation was then measured because the peak of ROS generation for both PMNLs and MNCs varied from one subject to another, whereas an increase occurred in all subjects. The AUC was calculated for NF- κ B binding activity, I κ B α , IKK α , IKK β , and p47^{phox} and then compared between the meal-fed and control subjects by using Student's t test for unpaired data. CRP and sICAM-1 concentrations were also normalized to a baseline value of 100% because the baseline concentrations varied markedly from one subjects to another. A P value < 0.05 was used to assess the significance for all statistical analyses. The results are presented as means \pm SEMs.

RESULTS

Glucose, lipid, and insulin concentrations

Plasma glucose concentrations did not change significantly after the meal challenge (Table 1). Plasma insulin concentration increased significantly at 1 h and remained elevated at 2 and 3 h when compared with baseline. Plasma triacylglycerol concentrations increased significantly 2 and 3 h after the meal challenge.

Plasma HDL, LDL, and cholesterol did not change significantly after food intake.

NF- κ B binding activity

NF- κ B binding to the consensus sequence oligonucleotide showed 2 bands on electrophoretic mobility shift assay. The upper band (I) was supershifted with the antibodies to both p65 and p50, whereas the second band (II) was supershifted with an antibody against p50 but not against p65 (Figure 1). NF- κ B (p65:p50 heterodimer) binding activity in MNC nuclear extracts increased significantly 1 and 2 h after the meal challenge and returned to baseline at 3 h ($P < 0.05$; Figure 1). The decrease in the p50:p50 complex (II) was not consistent in all the subjects. Only 4 subjects showed this decrease, whereas the other subjects showed no change in complex II formation.

I κ B α , IKK α , and IKK β concentrations

I κ B α protein concentrations, measured by Western blotting, decreased significantly 1 h after meal intake, as shown in Figure 2. This decrease persisted at 2 and 3 h ($P < 0.05$) when compared with baseline (0 h) and with the values for the control subjects (water intake). IKK α protein concentrations, measured by Western blotting, showed a significant increase 1, 2, and 3 h after meal intake when compared with both baseline and with the values for the control subjects ($P < 0.05$; Figure 3). IKK β protein concentrations did not increase after the meal challenge when com-




FIGURE 3. A: Representative Western blot showing the relative expression of inhibitor κ B kinase α (IKK α) in mononuclear cell homogenates 0, 1, 2, and 3 h after a 900-kcal mixed-meal challenge or water intake. B: Mean (\pm SEM) change in IKK α concentrations in mononuclear cells by densitometric quantitative analysis. There was a significant interaction between treatment (water compared with meal) and time ($P < 0.005$, two-factor ANOVA). *Significantly different from baseline (0 h), $P < 0.05$ (one-factor ANOVA for repeated measures followed by Dunnett's test). $n = 9$ for meal intake and $n = 8$ for water intake. C: Area under the curve (AUC) for IKK α protein expression after water and meal intakes. *Significantly different from water intake, $P < 0.05$ (Student's t test).

pared with baseline ($P = 0.28$; **Figure 4**). However, the AUC for IKK β was statistically significant after the meal challenge. This finding suggests that the increase in nuclear NF- κ B was due to the inhibition of expression of I κ B, which in turn was due to the induction of IKK α and IKK β , which phosphorylate I κ B α and cause its degradation.

p47^{phox} Subunit concentrations and ROS generation

The protein quantities of the p47^{phox} subunit of NADPH oxidase in MNC homogenates increased significantly at 1 h and remained elevated until 3 h when compared with both baseline and with the values for the control subjects ($P < 0.05$; **Figure 5**). All 9 subjects showed an increase in ROS generation by MNCs after meal intake. However, the time of the peak increase varied from one subject to another: in 1 subject the peak was at 1 h, in 5 subjects it was at 2 h, and in 3 subjects it was at 3 h. Baseline ROS generation by MNCs was 256 ± 140 mV (= 100%). Because of the interindividual variation in ROS generation, the AUC for all ROS generation was measured. The AUC for ROS generation by MNCs increased significantly (**Figure 6**; $P < 0.05$) after the mixed-meal challenge when compared with the AUC of ROS generation after the water challenge. Eight of 9 subjects showed an increase in ROS generation by PMNLs; the peak increase varied from one subject to another. Baseline ROS generation by PMNLs was 108 ± 55 mV. The AUC for ROS

generation by PMNLs also increased significantly (**Figure 6**; $P < 0.05$) after the mixed-meal challenge when compared with the AUC of ROS generation after the water challenge.

Plasma CRP, sICAM-1, and α -tocopherol concentrations

After the meal challenge, the increase in absolute plasma CRP concentrations was not significant. However, when values were normalized to a baseline concentration of 100% because of the interindividual variation in plasma CRP concentrations, this increase was significant ($P < 0.05$) when compared with one-factor ANOVA for the repeated measures followed by Dunn's test (**Table 2**). Water intake, on the other hand, did not cause any change in absolute or normalized plasma CRP concentrations. There was a significant interaction between treatment (water compared with the mixed meal) and time when CRP concentrations were compared between the 2 groups with two-factor ANOVA ($P < 0.05$). Plasma sICAM-1 concentrations did not change significantly after the meal challenge ($P = 0.119$) or water intake (**Table 2**). Plasma α -tocopherol concentrations did not change after the meal challenge (**Table 1**).

DISCUSSION

Our data show that after the intake of a 900-kcal mixed meal, there is a significant increase in intranuclear NF- κ B binding




FIGURE 4. A: Representative Western blot showing the relative expression of inhibitor κ B kinase β (IKK β) in mononuclear cell homogenates 0, 1, 2, and 3 h after a 900-kcal mixed-meal challenge or water intake. Note that IKK β concentrations did not increase significantly, largely because of variation in the magnitude of change after the meal. B: Mean (\pm SEM) change in IKK β concentrations in mononuclear cells by densitometric quantitative analysis. $n = 9$ for meal intake and $n = 8$ for water intake. There was no significant interaction between treatment (water compared with meal) and time (two-factor ANOVA). C: Area under the curve (AUC) for IKK β protein expression after water and meal intakes. *Significantly different from water intake, $P < 0.05$ (Student's t test).

activity by the proinflammatory p65:p50 heterodimer in MNCs. The increase was significant 1 and 2 h after the meal challenge and decreased 3 h after the meal challenge. This is important because the p65:p50 heterodimer is thought to be responsible for the transcription of proinflammatory genes, whereas the p50:p50 homodimer inhibits the activation of proinflammatory genes. Associated with this increase in NF- κ B binding activity in the nucleus was a decrease in cellular I κ B α . I κ B α is present in the cytosol, where it binds to NF- κ B and prevents its translocation into the nucleus (14, 15). It is noteworthy that proinflammatory stimuli such as endotoxin and cytokines (eg, tumor necrosis factor α) induce an increase in intranuclear NF- κ B and a decrease in I κ B (14–17). On the other hand, antiinflammatory agents cause an increase in I κ B α and a decrease in NF- κ B (18). The intake of 300 mL water did not cause a significant change in NF- κ B, I κ B, or any other index measured in the study.

The increase in IKK α and IKK β after the caloric challenge of the mixed meal is of interest because this enzyme is induced by endotoxin challenge and is responsible for the phosphorylation, ubiquitination, and eventual proteasomal degradation of I κ B α . This decrease in I κ B releases NF- κ B to translocate into the nucleus. There are recent data that show that IKK α may have an additional function: it may translocate into the nucleus and act as a coactivator of NF- κ B–induced tumor necrosis factor α and other proinflammatory gene expression (19).

p47^{phox} Subunit expression increased significantly at 1 h and persisted at an elevated concentration for 3 h. This suggests that

NADPH oxidase activity increased significantly after the meal, which is consistent with our previous data on the effects of glucose intake (8). Our data are thus consistent with the NF- κ B–mediated modulation of the p47^{phox} subunit. We previously showed that the suppression of intranuclear NF- κ B is associated with a decrease in p47^{phox} subunit expression, as observed after hydrocortisone (20), rosiglitazone (P Mohanty, A Aljada, H Ghanim et al, unpublished observations, 2001), troglitazone (21), and insulin (22, 23) injections, whereas an increase in NF- κ B is associated with a parallel increase in p47^{phox} subunit expression as observed after glucose intake (8). The essential role of NADPH oxidase and superoxide radical in the pathogenesis of atherosclerosis was recently shown. ApoE $-/-$ mice, in whom the p47^{phox} gene has been deleted, do not develop atherosclerotic lesions even when fed a high-cholesterol diet (24). Thus, it is of interest that p47^{phox} and superoxide radical get induced with each meal and that obesity is characterized by a marked increase in oxidative stress (4).

The magnitude of the peak increase in the p47^{phox} subunit after the intake of the mixed meal was similar to that observed after glucose intake, but the peak increase in ROS generation was at different times in the subjects studied. The lack of a significant increase in ROS generation at a single time point contrasts with the increases we previously showed after glucose (1), cream, and casein (8) challenges. This may be the result of the variability caused by the variable digestion of food before absorption and the peak of ROS generation being blunted by the innate antioxi-




FIGURE 5. A: Representative Western blot showing the relative expression of p47^{phox} subunit in mononuclear cell homogenates 0, 1, 2, and 3 h after a 900-kcal mixed-meal challenge or water intake. B: Mean (\pm SEM) change in p47^{phox} subunit protein expression in mononuclear cell homogenates by densitometric quantitative analysis. There was a significant interaction between treatment (water compared with meal) and time ($P < 0.05$, two-factor ANOVA). *Significantly different from baseline (0 h), $P < 0.05$ (one-factor ANOVA for repeated measures followed by Dunnett's test). $n = 9$ for meal intake and $n = 8$ for water intake. C: Area under the curve (AUC) for p47^{phox} subunit protein expression in mononuclear cell homogenates after water and meal intakes. *Significantly different from water intake, $P < 0.05$ (Student's t test).

dants in the mixed meal, including vitamins C and E (25). However, when the AUC of an increase from baseline in ROS generation at 1, 2, and 3 h was combined and comparisons were made




FIGURE 6. Mean (\pm SEM) reactive oxygen species (ROS) generation by mononuclear cells (MNCs) and polymorphonuclear leukocytes (PMNLs) after a 900-kcal mixed-meal challenge or water intake. *Significantly different from water intake, $P < 0.05$ (Student's t test for unpaired data). $n = 9$ for meal intake and $n = 8$ for water intake.

between the mixed-meal group and the control subjects, there was a significant increase in ROS generation by both MNCs and PMNLs in the mixed-meal group compared with the control subjects, who drank 300 mL water.

Our observations have relevance to the pathogenesis of atherosclerosis, in which inflammation may play a role through leukocyte-endothelial interactions. Such damage is of special relevance to diabetic patients because glycated proteins in these patients may trigger inflammation through advanced glycation end product receptors (26, 27). The proinflammatory changes induced by a meal may act in combination with the increase in oxidative stress, which is known to be increased in diabetes. Similarly, macrovascular disease in obesity may be related to increased oxidative stress, which we recently showed in obese subjects (4), and to the proinflammatory changes induced by meal intake. Thus, these observations may be relevant to the pathogenesis of atherosclerosis, which is a chronic inflammatory state of the arterial wall (28, 29), in both diabetes and obesity. It is also relevant to the reversal of atherosclerosis and the improvement in myocardial perfusion, which was shown after intense lifestyle changes by Ornish et al (30).

These observations may also help explain the abnormalities in brachial artery reactivity observed after consumption of the 900-kcal meal in the current study. Vogel et al (6) showed that such a meal results in a significant reduction in postischemic flow-

TABLE 2

Plasma C-reactive protein (CRP) and soluble intercellular adhesion molecule 1 (sICAM-1) concentrations after a 900-kcal mixed-meal challenge or water intake¹

	Baseline (0 h)	Time after meal challenge		
		1 h	2 h	3 h
Plasma CRP				
Meal (<i>n</i> = 9)				
Plasma CRP (ng/mL)	1444 (1189–2087) ²	1441 (1140–2021)	1546 (1224–2389)	1574 (1213–2443)
Normalized CRP (%)	100	104.0 ± 12.1	107. ± 16.0	128.0 ± 54.4 ³
Water (<i>n</i> = 8)				
Plasma CRP (ng/mL)	1085 (521–1502)	1085 (531–1549)	1204 (533–1604)	1063 (592–1477)
Normalized CRP (%)	100	100.4 ± 9.1	104.4 ± 8.6	104.9 ± 16.9
Plasma sICAM-1				
Meal (<i>n</i> = 9)				
Plasma sICAM-1 (ng/mL)	231.5 ± 19.7	244.5 ± 18.6	260.9 ± 21.2	242.6 ± 19.6
Normalized sICAM-1 (%)	100	106.9 ± 11.5	113.5 ± 13.0	106.1 ± 17.2
Water (<i>n</i> = 8)				
Plasma sICAM-1 (ng/mL)	170.0 ± 9.3	167.3 ± 8.0	174.1 ± 10.6	175.2 ± 7.1
Normalized sICAM-1 (%)	100	99.0 ± 7.8	102.3 ± 4.3	104.3 ± 13.1

¹ There was a significant interaction between treatment (water compared with meal) and time when CRP concentrations were compared between the 2 groups with two-factor ANOVA ($P < 0.05$). sICAM-1 did not change significantly after the meal or water intake.

² Median; interquartile range in parentheses (all such values).

³ Significantly different from baseline, $P < 0.05$ (one-factor repeated-measures ANOVA followed by Dunnett's test).


mediated vasodilation of the brachial artery. Furthermore, they showed that prior intake of vitamins E and C prevents this abnormality (31). They concluded that the abnormality in vascular reactivity induced by the meal was mediated through an increase in ROS generation and a possible reduction in the bioavailability of endothelial nitric oxide. Our observations provide actual evidence that confirms their hypothesis in terms of increased ROS generation and altered endothelium-mediated vasodilatation probably through reduced nitric oxide bioavailability.

The concomitant increase in IKK α and IKK β expression, decrease in I κ B α , and increase in intranuclear NF- κ B are consistent with leukocytic activation and a proinflammatory effect of the intake of a moderate-sized meal. It is of interest that these processes get triggered within 1 h of a meal and remain active for ≥ 3 h and that CRP also increases after such a meal. The duration of this proinflammatory effect is relevant because the time for another meal is usually 4–5 h after a meal, with the potential for further NF- κ B activation, ROS load, and potential proinflammatory changes. It is possible that chronic overeating may result in permanent increases in NF- κ B binding activity and in total expression as a protein in the cell. This area needs further investigation because the state of obesity is known to be associated with an increase in oxidative stress and an increase in plasma concentrations of proinflammatory mediators such as tumor necrosis factor α , interleukin 6, and CRP (32–34).

The fact that insulin is secreted in response to the meal raises the theoretical possibility that it may mediate this proinflammatory effect. However, we already showed that insulin has a potent and rapid antiinflammatory effect both in vivo (22, 23) and in vitro (35, 36). Insulin suppresses intranuclear NF- κ B, induces I κ B α , and suppresses the p47^{phox} subunit of NADPH oxidase, ICAM-1, MCP-1, and CRP, all of which are proinflammatory.

Our data are relevant to the concept that, whereas insulin and insulin sensitizers of the thiazolidinediones class have an ROS-suppressive and a profound antiinflammatory effect, macronutrient intake induces oxidative stress and proinflammatory

changes. Thus, our current understanding of the relation between macronutrient intake and insulin, which contains metabolic implications only, has to incorporate the respective inflammatory and antiinflammatory effects of the macronutrient intake and insulin action.

In conclusion, a moderate-sized mixed meal results in a significant increase in the proinflammatory transcription factor NF- κ B in the nucleus, a decrease in I κ B α (which is probably a consequence of the increase in IKK α and IKK β), and a parallel increase in the cellular p47^{phox} subunit, which is the key protein component of NADPH oxidase and in ROS generation. These changes occur within 1 h of meal consumption and last for ≥ 3 h. These observations are consistent with potent inflammatory changes and raise fundamental issues about the relation between food intake, oxidative damage, inflammation, and atherosclerosis and about the ideal way to eat and modify lifestyle. 

REFERENCES

- Mohanty P, Hamouda W, Garg R, Aljada A, Ghanim H, Dandona P. Glucose challenge stimulates reactive oxygen species (ROS) generation by leukocytes. *J Clin Endocrinol Metab* 2000;85:2970–3.
- Mohanty P, Ghanim H, Hamouda W, Aljada A, Garg R, Dandona P. Both lipid and protein intakes stimulate increased generation of reactive oxygen species by polymorphonuclear leukocytes and mononuclear cells. *Am J Clin Nutr* 2002;75:767–72.
- Dandona P, Mohanty P, Hamouda W, et al. Inhibitory effect of a two day fast on reactive oxygen species (ROS) generation by leukocytes and plasma ortho-tyrosine and meta-tyrosine concentrations. *J Clin Endocrinol Metab* 2001;86:2899–902.
- Dandona P, Mohanty P, Ghanim H, et al. The suppressive effect of dietary restriction and weight loss in the obese on the generation of reactive oxygen species by leukocytes, lipid peroxidation and protein carbonylation. *J Clin Endocrinol Metab* 2001;86:355–62.
- Raitakari OT, Lai N, Griffiths K, McCredie R, Sullivan D, Celermajer DS. Enhanced peripheral vasodilation in humans after a fatty meal. *J Am Coll Cardiol* 2000;36:417–22.
- Vogel RA, Corretti MC, Plotnick GD. Effect of a single high-fat meal on endothelial function in healthy subjects. *Am J Cardiol* 1997;79:350–4.
- Ceriello A, Bortolotti N, Motz E, et al. Meal-induced oxidative stress and

- low-density lipoprotein oxidation in diabetes: the possible role of hyperglycemia. *Metabolism* 1999;48:1503–8.
8. Aljada A, Ghanim H, Mohanty P, Assian E, Dandona P. Glucose intake stimulates intranuclear NFkB and p47phox in mononuclear cells. *J Clin Endocrinol Metab* (in press).
 9. Andrews NC, Faller DV. A rapid micropreparation technique for extraction of DNA-binding proteins from limiting numbers of mammalian cells. *Nucleic Acid Res* 1991;19:2499.
 10. Dandona P, Karne R, Ghanim H, Hamouda W, Aljada A, Magsino CH Jr. Carvedilol inhibits reactive oxygen species generation by leukocytes and oxidative damage to amino acids. *Circulation* 2000;101:122–4.
 11. Dandona P, Thusu K, Hafeez R, Abdel-Rahman E, Chaudhuri A. Effect of hydrocortisone on oxygen free radical generation by mononuclear cells. *Metabolism* 1998;47:788–91.
 12. Tosi MF, Hamedani A. A rapid, specific assay for superoxide release from phagocytes in small volumes of whole blood. *Am J Clin Pathol* 1992;97:566–73.
 13. Hancock JT, Jones OT. The inhibition by diphenyleneiodonium and its analogues of superoxide generation by macrophages. *Biochem Genet* 1987;242:103–7.
 14. Baeuerle PA, Henkel T. Function and activation of NF-kappa B in the immune system. *Annu Rev Immunol* 1994;12:141–79.
 15. Barnes PJ, Karin M. Nuclear factor-kappaB: a pivotal transcription factor in chronic inflammatory diseases. *N Engl J Med* 1997;336:1066–71.
 16. Zuckerman SH, Evans GF. Endotoxin tolerance: in vivo regulation of tumor necrosis factor and interleukin-1 synthesis is at the transcriptional level. *Cell Immunol* 1992;140:513–9.
 17. Manning AM, Bell FP, Rosenbloom CL, et al. NF-kappa B is activated during acute inflammation in vivo in association with elevated endothelial cell adhesion molecule gene expression and leukocyte recruitment. *J Inflamm* 1995;45:283–96.
 18. Aljada A, Ghanim H, Assian E, et al. Increased IkappaB expression and diminished nuclear NF-kappaB in human mononuclear cells following hydrocortisone injection. *J Clin Endocrinol Metab* 1999;84:3386–9.
 19. Israel A. Signal transduction: a regulator branches out. *Nature* 2003;423:596–7.
 20. Dandona P, Aljada A, Ghanim H, Mohanty P, Hamouda W, Al-Haddad W. Acute suppressive effect of hydrocortisone on p47 subunit of nicotinamide adenine dinucleotide phosphate oxidase. *Metabolism* 2001;50:548–52.
 21. Ghanim H, Garg R, Aljada A, et al. Suppression of nuclear factor-kappaB and stimulation of inhibitor kappaB by troglitazone: evidence for an anti-inflammatory effect and a potential antiatherosclerotic effect in the obese. *J Clin Endocrinol Metab* 2001;86:1306–12.
 22. Dandona P, Aljada A, Mohanty P, et al. Insulin inhibits intranuclear nuclear factor kappaB and stimulates IkappaB in mononuclear cells in obese subjects: evidence for an anti-inflammatory effect? *J Clin Endocrinol Metab* 2001;86:3257–65.
 23. Dandona P, Aljada A, Mohanty P. The anti-inflammatory and potential anti-atherogenic effect of insulin: a new paradigm. *Diabetologia* 2002;45:924–30.
 24. Patricia A, Barry-Lane, Patterson C, Holland S, Yeh E, Runge M. p47phox is required for atherosclerotic lesion progression in AopE^{-/-} mice. *J Clin Invest* 2001;108:1513–22.
 25. Lentner C, ed. Units of measurement, body fluids, composition of the body, nutrition. West Caldwell, NJ: Ciba-Geigy Corp, 1981.
 26. Schmidt AM, Hofmann M, Taguchi A, Yan SD, Stern DM. RAGE: a multiligand receptor contributing to the cellular response in diabetic vasculopathy and inflammation. *Semin Thromb Haemost* 2000;26:485–93.
 27. Schmidt AM, Stern DM. RAGE: a new target for the prevention and treatment of the vascular and inflammatory complications of diabetes. *Trends Endocrinol Metab* 2000;11:368–75.
 28. Ross R. Atherosclerosis—an inflammatory disease. *N Engl J Med* 1999;340:115–26.
 29. Schwartz CJ, Valente AJ, Sprague EA, Kelley JL, Suenram CA, Rozek MM. Atherosclerosis as an inflammatory process. The roles of the monocyte-macrophage. *Ann N Y Acad Sci* 1985;454:115–20.
 30. Ornish D, Scherwitz LW, Billings JH, et al. Intensive lifestyle changes for reversal of coronary heart disease. *JAMA* 1998;280:2001–7.
 31. Plotnick GD, Corretti MC, Vogel RA. Effect of antioxidant vitamins on the transient impairment of endothelium-dependent brachial artery vasoactivity following a single high-fat meal. *JAMA* 1997;278:1682–6.
 32. Dandona P, Weinstock R, Thusu K, Abdel-Rahman E, Aljada A, Wadden T. Tumor necrosis factor-alpha in sera of obese patients: fall with weight loss. *J Clin Endocrinol Metab* 1998;83:2907–10.
 33. Yudkin JS, Stehouwer CD, Emeis JJ, Coppack SW. C-reactive protein in healthy subjects: associations with obesity, insulin resistance, and endothelial dysfunction: a potential role for cytokines originating from adipose tissue? *Arterioscler Thromb Vasc Biol* 1999;19:972–8.
 34. Pradhan AD, Manson JE, Rifai N, Buring JE, Ridker PM. C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *JAMA* 2001;286:327–34.
 35. Aljada A, Ghanim H, Saadeh R, Dandona P. Insulin inhibits NFkappaB and MCP-1 expression in human aortic endothelial cells. *J Clin Endocrinol Metab* 2001;86:450–3.
 36. Aljada A, Saadeh R, Assian E, Ghanim H, Dandona P. Insulin inhibits the expression of intercellular adhesion molecule-1 by human aortic endothelial cells through stimulation of nitric oxide. *J Clin Endocrinol Metab* 2000;85:2572–5.

