

AIDS:

20 August 1998 - Volume 12 - Issue 12 - p 1553-1555

Correspondence

Serum antibodies to dietary antigens in patients with HIV-1 infection

Wahnschaffe, U¹; Zippel, T¹; Schmidt, W¹; Schneider, T²;
Schulzke, J-D¹; Riecken, E-O¹; Ullrich, R¹



Author Information

¹Medical Clinic, Department of Gastroenterology and Infectious Diseases, Universitätsklinikum Benjamin Franklin, Freie Universität Berlin, Berlin

²Medical Clinic, Department of Medicine, University of the Saarland, Homburg/Saar, Germany.

Date of receipt: 5 March 1997; accepted: 11 March 1998.

Diarrhoea is a common complication of HIV-1 infection [1], but in a considerable proportion of patients, especially in earlier stages of the disease, the cause of diarrhoea remains unclear despite extensive investigation [2]. Increased levels of serum antibodies against food proteins have been described in HIV-infected children [3], similar to patients with non-IgE-mediated gastrointestinal food hypersensitivity [4], coeliac disease [5], or chronic inflammatory bowel disease [6]. These abnormalities are thought to result from increased gut permeability leading to increased uptake of dietary antigens and an aberrant mucosal IgG response in these patients [7]. Increased intestinal permeability [8] and an increase of IgG in mucosal secretions [9] have also been found in patients infected with HIV-1. Since abnormal immunity to dietary antigens may contribute to the pathogenesis of diarrhoea, we investigated serum antibodies to food proteins in HIV-1-infected patients with and without diarrhoea.

Sera from 70 consecutive HIV-infected patients (65 men, five women) attending the HIV outpatient clinic were studied. Forty-three patients had AIDS. The median age was 44 years (range, 29-64 years) and the median CD4 T-cell count was $65 \times 10^6/l$ (range, $4-987 \times 10^6/l$). Eleven HIV-infected patients had diarrhoea. In six of these patients an enteric pathogen was found (*Salmonella* sp. and microsporidia infection each in two patients, *Mycobacterium avium* complex and *Isospora belli* infection in one patient, and one patient with microsporidia had additional coronavirus infection). Blood samples from 20 healthy controls (15 men, five women; mean age, 55.5 years; range, 18-73 years) not at risk for HIV infection and without evidence for immune defect served as controls.

Serum IgG, IgA and IgM titres against the dietary antigens β -lactoglobulin, α -lactalbumin, ovalbumin, and soy protein, and the non-dietary antigen tetanus toxoid were measured by enzyme-linked immunosorbent assay. Because serum immunoglobulins are unspecifically increased in HIV-infected patients compared with

controls, antibody titres were corrected for serum immunoglobulin concentrations as measured by single radial immunodiffusion. Serum IgG, IgA and IgM antibody levels against all food proteins (except anti- β -lactoglobulin IgM) were significantly higher in HIV-infected patients than in controls. In contrast, IgG antibody titres against tetanus toxoid were decreased in HIV-infected patients compared with controls (each $P < 0.05$; Fig. 1). No differences were found for IgA and IgM antibody titres against tetanus toxoid. When antibody titres were expressed in relation to the respective serum immunoglobulin concentration, IgG antibody levels against all food proteins tested were still increased (each $P < 0.01$), although IgG against tetanus toxoid was decreased in HIV-infected patients compared with controls. Relative IgA and IgM titres against the tested proteins were similar in HIV-infected patients and controls. No differences were found between HIV-infected patients with and without AIDS. Median antibody titres for IgG, IgA, and IgM against all tested food proteins were higher in HIV-infected patients with diarrhoea than in patients who had no diarrhoea, but a significant difference was found only for IgG antibodies against ovalbumin ($P < 0.05$).




Fig. 1

Our results show an increase of serum antibodies against four common food proteins in HIV-infected patients. This is not due to unspecific polyclonal B-cell activation because immunoglobulin levels were still increased when antibody levels were corrected for total immunoglobulin concentration. In addition, no increase of serum antibodies against the non-dietary protein tetanus toxoid was observed. The appearance of antibodies to food proteins in serum indicates an abnormal access of these antigens to the mucosal immune system, probably due to an impaired epithelial barrier function [8]. Although we did not specifically investigate local humoral immunity, the increased mucosal production of IgG in HIV-infected patients [10] and the high IgG levels in duodenal secretions of HIV-infected patients [11] indicate considerable transudation of serum antibodies into the gut. In contrast to the immune exclusion mediated by secretory IgA, IgG antibodies to food proteins will form immune complexes if their respective antigens are present in the mucosa and will activate complement [12]. The resulting mucosal inflammation that is frequently observed in the intestine of HIV-infected patients [13] may not only establish a vicious circle perpetuating impairment of epithelial barrier function but could also contribute to the pathogenesis of diarrhoea. The increased immunity to food proteins in HIV infection shown in our study suggests that HIV-infected patients might benefit from nutritional interventions. This should be tested primarily in HIV-infected patients with high titres of serum antibodies to dietary antigens.

References

1. Ullrich R, Heise W, Bergs C, L'age M, Riecken EO, Zeitz M: **Gastrointestinal symptoms in patients infected with human immunodeficiency virus:**

relevance of infective agents isolated from gastrointestinal tract. *Gut* 1992, **33**:1080-1084.

2. Wilcox M, Schwartz DA, Cotsonis G, Thompson III SE: **Chronic unexplained diarrhea in human immunodeficiency virus infection: determination of the best diagnostic approach.** *Gastroenterology* 1996, **110**:30-37.

3. Quesnel A, Moja PH, Blanche S, Griscelli C, Genin C: **Early impairment of gut mucosal immunity in HIV-1-infected children.** *Clin Exp Immunol* 1994, **97**:380-385.

4. Pearson JR, Kingston D, Shiner M: **Antibody production to milk proteins in the jejunal mucosa of children with cow's milk protein intolerance.** *Pediatr Res* 1983, **17**:406-412.

5. Hvatum M, Brandtzaeg P: **Serum IgG subclass antibodies to a variety of food antigens in patients with coeliac disease.** *Gut* 1992, **33**:632-638.

6. Knoflauch P, Park BH, Cunningham R, Weiser MM, Albin B: **Serum antibodies to cow's milk proteins in ulcerative colitis and Crohn's disease.** *Gastroenterology* 1987, **92**:479-485.

7. Husby S, Foged N, Host A, Svehag SE: **Passage of dietary antigens into the blood of children with coeliac disease. Quantification and size distribution of absorbed antigens.** *Gut* 1987, **28**:1062-1072.

8. Stockmann M, Fromm M, Schmitz H, Schmidt W, Riecken E-O, Schulzke J-D: **Duodenal biopsies of HIV-infected patients with diarrhoea exhibit epithelial barrier defects but no active secretion.** *AIDS* 1998, **12**:43-51.

9. Schneider T, Zippel T, Schmidt W, *et al.*: **Abnormal predominance of IgG in HIV-specific antibodies produced by short-term cultured duodenal biopsy specimens from HIV-infected patients.** *J Acquir Immune Defic Syndr Hum Retrovirol* 1998, **16**:333-339.

10. Schneider T, Zippel T, Schmidt W, *et al.*: **Increased immunoglobulin G production by short term cultured duodenal biopsy samples from HIV infected patients.** *Gut* 1998, **42**:357-361.

11. Janoff ED, Jackson S, Wahl SM, Thomas K, Peterman JH, Smith PD: **Intestinal mucosal immunoglobulins during human immunodeficiency virus type 1 infection.** *J Infect Dis* 1994, **170**:299-307.

12. Halstensen TS, Das KM, Bandtzaeg P: **Epithelial deposits of immunoglobulin G1 and activated complement colocalise with the Mr 40 kD putative autoantigen in ulcerative colitis.** *Gut* 1993, **34**:650-657.

13. Kotler DP, Reka S, Cayton F: **Intestinal mucosal inflammation associated with human immunodeficiency virus infection.** *Dig Dis Sci* 1993, **38**:1119-1127.

© 1998 Lippincott Williams & Wilkins, Inc.